

Refraction and absorption of x rays by laser-dressed atoms

Christian Buth, Robin Santra, and Linda Young
Argonne National Laboratory, Argonne, Illinois 60439, USA
(Dated: 18 March 2008)

X-ray refraction and absorption by neon atoms under the influence of an 800 nm laser with an intensity of 10^{13} W/cm² is investigated. For this purpose, we use an *ab initio* theory suitable for optical strong-field problems. Its results are interpreted in terms of a three-level model. On the Ne $1s \rightarrow 3p$ resonance, we find electromagnetically induced transparency (EIT) for x rays. Our work opens novel perspectives for ultrafast x-ray pulse shaping.

PACS numbers: 32.30.Rj, 32.80.Fb, 32.80.Rm

We consider atoms in the light of a linearly polarized laser which are probed by linearly polarized x rays [Fig. 1]. Such a setting is referred to as a two-color problem and has proved to be a very beneficial to probe strong-field physics and to manipulated x rays [1, 2, 3, 4, 5, 6, 7].

We assume a Ti:sapphire laser at 800 nm with 10^{13} W/cm². The atomic ground-state electrons remain essentially unperturbed in the laser light; they are neither excited nor ionized. However, laser dressing of the core-excited states introduces strong-field physics because, for the laser parameters employed here, the Keldysh parameter [8] is $\gamma = \sqrt{I_{1s^{-1}3p}/(2U_p)} = 1.5 \approx 1$. The ionization potential of a core-excited state with the electron in the $3p$ Rydberg orbital is denoted by $I_{1s^{-1}3p}$ and the ponderomotive potential of the laser is U_p [3].

For $\gamma \approx 1$, neither perturbation theory nor a tunneling description is adequate. Therefore, we have developed an *ab initio* theory described in Ref. 1. The electronic structure of the atom is described in the Hartree-Fock-Slater approximation where a complex absorbing potential (CAP) is used to treat continuum electrons. Nonrelativistic quantum electrodynamics is used to treat the atom in the two-color light. The interaction with the laser is described in terms of a Floquet-type matrix whereas the x-ray probe is treated in second-order (one-photon) perturbation theory [9]. For core-excited states, we need to take into account the Auger decay rate of a K -shell hole in neon $\Gamma_{1s} = 0.27$ eV. The absorption cross section for x rays with photon energy ω_X is found




FIG. 1: (Color online) Laser-dressed neon atom probed by x rays.

to be [1]

$$\sigma_{1s}(\omega_X, \vartheta_{LX}) = \sigma_{1s}^{\parallel}(\omega_X) \cos^2 \vartheta_{LX} + \sigma_{1s}^{\perp}(\omega_X) \sin^2 \vartheta_{LX} . \quad (1)$$

where ϑ_{LX} denotes the angle between the laser and x-ray polarization vectors. The calculations were carried out with the DREYD and PULSEPROP programs [10]. We use the computational parameters from Refs. [1, 3]. For the index of refraction [Fig. 4], we need 400 radial wavefunctions.




FIG. 2: (Color online) X-ray photoabsorption cross section of neon near the K edge with laser dressing and without it.

The calculated x-ray photoabsorption cross section of a neon atom is displayed in Fig. 2 with laser dressing and without. The prominent absorption feature in the top panel at 867.4 eV arises due to the $1s \rightarrow 3p$ resonance. The weaker peaks are associated with $1s \rightarrow np$ transitions with $n \geq 4$. The K edge lies at 870.2 eV. The most eye-catching impact of the laser dressing can be seen in the vicinity of the $1s \rightarrow 3p$ resonance, which becomes transparent for $\vartheta_{LX} = 0^\circ$ [3].

To understand the influence of the laser, we make a Λ -type model for neon shown in Fig. 3. We use the states $1s^{-1}3p$, $1s^{-1}3s$, and the atomic ground state. The laser photon energy is ω_L whereas $\Gamma_{\text{Ne } 1s^{-1}3p}$ and




FIG. 3: (Color online) Λ -type three-level model for neon.

$\Gamma_{\text{Ne } 1s^{-1} 3s}$ denote the decay widths of the respective core-excited states. We find the overall agreement of the model curves with the *ab initio* data to be reasonably well. The model explains the transparency induced by the laser in the middle panel of Fig. 3 in terms of a splitting of the $1s^{-1} 3p$ and $1s^{-1} 3s$ states into an Autler-Townes doublet. Since the laser causes a suppression of resonant absorption, we call this electromagnetically induced transparency (EIT) for x rays [3]. Dipole selection rules dictate that the laser can couple only $1s^{-1} 3p_z$ to $1s^{-1} 3s$ in a one-photon processes; so only for $\vartheta_{\text{LX}} = 0^\circ$, EIT occurs. The suppression for $\vartheta_{\text{LX}} = 90^\circ$ is a consequence of laser-induced line broadening.




FIG. 4: (Color online) Refractive index of neon without laser dressing.

The refractive index is a classical quantity from Maxwell's equations. In Fig. 4, we show the real part $n(\omega_X)$ for neon in the x-ray regime without laser dressing at normal temperature and pressure [7]. It follows from $n(\omega_X) = 1 + 2\pi n_{\#} \alpha_I(\omega_X)$, where the number density of neon atoms is $n_{\#}$. Further, $\alpha_I(\omega_X)$ is the dynamic polarizability which we compute using $\hat{H}_{\text{I,X}} = \alpha \hat{\vec{p}} \cdot \vec{A}_X + \frac{1}{2} \alpha^2 \vec{A}_X^2$ (in atomic units) for the interaction Hamilton with the x rays in dipole approximation [7]. The momentum operator is $\hat{\vec{p}}$ and the vector potential is $\vec{A}_X$. We compare our theoretical results with the data from Fig. 4 of Ref. 11 which was computed from experimental cross section data. The agreement is generally good; only the feature on the $1s \rightarrow 3p$ resonance is not reproduced by the semi-experimental data due to limited




FIG. 5: (Color online) Dynamic polarizability of neon near the K edge.

resolution.

The impact of the laser dressing on neon is investigated for the dynamic polarizability; it is plotted in Fig. 5 for the three cases considered in Fig. 2. In this paragraph, we follow Ref. 3 and use $\hat{H}_X = -\vec{r} \cdot \vec{E}_X$ in dipole approximation for the interaction with the x rays. The electron position is $\vec{r}$ and the electric field of the x rays is $\vec{E}_X$. The overall polarizability and its change is small for x rays. Laser dressing causes a suppression near the resonance in contrast to the optical domain. The Λ -type model [Fig. 3] reproduces the structure of $\alpha_I(\omega_X, \vartheta_{\text{LX}})$ [3].




FIG. 6: (Color online) EIT-based generation of two ultrashort x-ray pulses with well-defined time delay.

EIT for x rays opens up an exciting novel prospect: it can be used to imprint pulse shapes of the optical dressing laser onto a comparatively long x-ray pulse. Yet only amplitude modulation is practically feasible due to the small refraction and dispersion in the x-ray domain [Figs. 4 and 5]. In Fig. 6, the propagation of two-color light through

a neon gas cell is shown. First, the x-ray and laser pulses are still outside the gas cell. Second, the first of the two laser pulses overlaps with the x-ray pulse inside the gas cell, thereby substantially enhancing x-ray transmission in comparison to the laser-off case. Third, after propagation through the gas cell, two ultrashort x-ray pulses emerge. This EIT-based pulse shaping opens up a route to ultrafast all x-ray pump-probe experiments [3].

C.B.'s research was partly funded by a Feodor Lynen Research Fellowship from the Alexander von Humboldt Foundation. C.B.'s, R.S.'s, and L.Y.'s work was supported by the Office of Basic Energy Sciences, Office of Science, U.S. Department of Energy, under Contract No. DE-AC02-06CH11357.

-
- [1] C. Buth and R. Santra, Phys. Rev. A **75**, 033412 (2007), arXiv:physics/0611122.
 - [2] Z.-H. Loh, M. Khalil, R. E. Correa, R. Santra, C. Buth, and S. R. Leone, Phys. Rev. Lett. **98**, 143601 (2007), arXiv:physics/0703149.

- [3] C. Buth, R. Santra, and L. Young, Phys. Rev. Lett. **98**, 253001 (2007), arXiv:0705.3615.
- [4] R. Santra, C. Buth, E. R. Peterson, R. W. Dunford, E. P. Kanter, B. Krässig, S. H. Southworth, and L. Young, J. Phys.: Conf. Ser. **88**, 012052 (2007), arXiv:0712.2556.
- [5] C. Buth and R. Santra, Phys. Rev. A **77**, 013413 (2008), arXiv:0711.3203.
- [6] E. R. Peterson, C. Buth, D. A. Arms, R. W. Dunford, E. P. Kanter, B. Krässig, E. C. Landahl, S. T. Pratt, R. Santra, S. H. Southworth, et al., Appl. Phys. Lett. **92**, 094106 (2008), arXiv:0802.1894.
- [7] C. Buth and R. Santra, manuscript in preparation (2008).
- [8] L. Keldysh, Sov. Phys. JETP **20**, 1307 (1965).
- [9] C. Buth, R. Santra, and L. S. Cederbaum, Phys. Rev. A **69**, 032505 (2004), arXiv:physics/0401081.
- [10] C. Buth and R. Santra, *FELLA – the free electron laser atomic, molecular, and optical physics program package*, Argonne National Laboratory, Argonne, Illinois, USA (2008), version 1.3.0, with contributions by Mark Baertschy, Kevin Christ, Chris H. Greene, Hans-Dieter Meyer, and Thomas Sommerfeld.
- [11] G. Liggett and J. S. Levinger, J. Opt. Soc. Am. **58**, 109 (1968).